



Arklow Bank Wind Park 2

Environmental Impact Assessment Report

Volume III, Appendix 12.4: Offshore Ornithology Technical Report -
Collision Risk Model Input Parameters



MacArthur
Green

Arklow Bank Wind Park 2

**Technical Appendix 12.04
Offshore Ornithology**

**Collision Risk Model Input
Parameters**

Date: 2 May 2024

Tel: 0141 342 5404

Web: www.macarthurgreen.com

Address: 93 South Woodside Road | Glasgow | G20 6NT

Document Quality Record

Version	Date	Status	Author	Reviewed by	Approved by
1.0	02/05/2024	Final (External)	MacArthur Green	GoBe Consultants	Sure Partners Limited

MacArthur Green is helping to combat the climate crisis through working within a carbon negative business model. Read more at www.macarthurgreen.com.



CONTENTS

Glossary	III
Acronyms.....	III
Units	IV
1 OFFSHORE ORNITHOLOGY TECHNICAL REPORT: COLLISION RISK MODEL INPUT PARAMETERS.....	1
1.1 Introduction	1
1.2 Collision Risk Model Input Parameters.....	1
1.2.1 Avoidance Rate.....	1
1.3 References.....	5

LIST OF TABLES

Table 12.4.1: Species biometrics used in the CRM. Mean values and standard deviations (where applicable).....	2
Table 12.4.2: Monthly mean densities (and 95% confidence intervals) of birds in flight within the Array Area used in the deterministic CRM.....	3
Table 12.4.3: Windfarm and turbine specifications used in the CRM.	4
Table 12.4.4: Wind turbine Monthly percentage operational time used in the CRM.	4

Glossary

Term	Meaning
Arklow Bank Wind Park 2 – Offshore Infrastructure	“The Proposed Development”, Arklow Bank Wind Park 2 Offshore Infrastructure: This includes all elements under the existing Maritime Area Consent.
Arklow Bank Wind Park 2 (ABWP2) (The Project)	<p>Arklow Bank Wind Park 2 (ABWP2) (The Project) is the onshore and offshore infrastructure. This EIAR is being prepared for the Offshore Infrastructure. Consents for the Onshore Grid Infrastructure (Planning Reference 310090) and Operations Maintenance Facility (Planning Reference 211316) has been granted on 26th May 2022 and 20th July 2022, respectively.</p> <ul style="list-style-type: none"> Arklow Bank Wind Park 2 Offshore Infrastructure: This includes all elements to be consented in accordance with the Maritime Area Consent. This is the subject of this EIAR and will be referred to as ‘the Proposed Development’ in the EIAR. Arklow Bank Wind Park 2 Onshore Grid Infrastructure: This relates to the onshore grid infrastructure for which planning permission has been granted. Arklow Bank Wind Park 2 Operations and Maintenance Facility (OMF): This includes the onshore and nearshore infrastructure at the OMF, for which planning permission has been granted. Arklow Bank Wind Park 2 EirGrid Upgrade Works: any non-contestable grid upgrade works, consent to be sought and works to be completed by EirGrid.
Array Area	The Array Area is the area within which the Wind Turbine Generators (WTGs), the Offshore Substation Platforms (OSPs), and associated cables (export, inter- array and interconnector cabling) and foundations will be installed.
Nocturnal Activity Factor	<p>Nocturnal Activity Factors indicate the amount of flight activity at night as a proportion of daytime flight activity.</p> <p>These factors were derived from reviews of seabird activity reported in Garthe and Hüppop (2004) which ranked species from 1 to 5 (1 low, 5 high) for relative nocturnal activity. These rates were subsequently modified for the purposes of CRM into 1 = 0%, 2 = 25%, 3 = 50%, 4 = 75% and 5 = 100% flying activity at night.</p> <p>For example, a nocturnal activity factor of 2 assumes that on average, nocturnal activity is around 25% of daytime level.</p>

Acronyms

Term	Meaning
CRM	Collision Risk Model/Modelling
EIA	Environmental Impact Assessment
JNCC	Joint Nature Conservation Committee
LAT	Lowest Astronomical Tide
MHW	Mean High Water
MSL	Mean Sea Level
NAF	Nocturnal Activity Factor

Term	Meaning
RPM	Revolutions Per Minute
SNCB	Statutory Nature Conservation Bodies
WTG	Wind Turbine Generator
UK	United Kingdom

Units

Unit	Description
Birds/km ²	Birds per square kilometre (density)
m	Metre (distance)
m/s	Metres per second (speed)

1 OFFSHORE ORNITHOLOGY TECHNICAL REPORT: COLLISION RISK MODEL INPUT PARAMETERS

1.1 Introduction

1. This Technical Report provides tables of the input parameters used in the deterministic Collision Risk Modelling (CRM) for the Array Area to inform the Environmental Impact Assessment (EIA) of the Arklow Bank Wind Park 2 Offshore Infrastructure (hereafter referred to as ‘the Proposed Development’). Collisions were estimated using the stochastic Band (2012) CRM, option 2 (generic flight heights).
2. Species were selected for inclusion in the CRM on the basis of a combination of factors, including the estimated density of birds recorded in-flight and knowledge of the species typical flight height range. For example, species including guillemot, razorbill and Manx shearwater fly close to the sea surface, well below the lower rotor tip heights, and therefore even at high densities the risk of collision is considered to be negligible. Therefore, these and similar low-flying species have been scoped out of the collision assessment (Johnson *et al.*, 2014). However, all other seabird species, even those recorded in very low densities, were included in the assessment to ensure that all potential impacts were considered.

1.2 Collision Risk Model Input Parameters

3. CRM input parameters are provided in the following tables:
 - a) Table 12.4.1: density (birds/km²) of birds in flight within the Array Area in each month, presented as the mean and upper and lower 95% confidence range derived from 1,000 nonparametric bootstrap simulations (the monthly values were derived as the mean of two months of survey density data and the 95% confidence intervals are derived from bootstrap data);
 - b) Table 12.4.2: biometrics for each species modelled (e.g. wingspan, body length, etc.), nocturnal activity factors and avoidance rates; and
 - c) Table 12.4.3 and Table 12.4.4: the wind farm and wind turbine operating parameters. Note that three turbine models are under consideration.
4. Generic flight height data were used as provided by Johnston *et al.*, (2014).

1.2.1 Avoidance Rate

5. A key parameter in the CRM is the avoidance rate, which accounts for the fact that birds will actively avoid colliding with the rotors (at a range of scales), while the baseline survey data are collected before turbines are installed. United Kingdom (UK) Statutory Nature Conservation Bodies (SNCBs) produced guidance on the rates to use for key collision risk species (Joint Nature Conservation Committee (JNCC) *et al.*, 2014) following a review conducted for Marine Scotland (Cook *et al.* 2014). Following this a joint industry study was conducted in an operational windfarm, using a combination of direct observations, radar and cameras (Skov *et al.*, 2018), which generated revised (higher) avoidance rates for gannet and kittiwake. Bowgen and Cook (2018)

were commissioned by the UK SNCBs to conduct an independent analysis using the same data and generated slightly more conservative avoidance rates.

6. More recently the University of Exeter was commissioned by JNCC to review and update seabird avoidance rates (Ozsanlav-Harris, *et al.*, 2023), which also recommended increases in avoidance for most species. Following this the avoidance rates recommended by Natural England are:

- Gannet 99.76% (inc. 70% macro avoidance; NB macro avoidance refers to avoidance of the windfarm as a whole – see Natural England 2022 for further explanation),
- Kittiwake 99.3%;
- Herring gull, lesser black-backed gull, great black-backed gull 99.4%;
- Little gull, common gull, black-headed gull 99.5%; and
- All other species 99.1%.

7. The remaining parameters are provided in Table 12.4.1.

Table 12.4.1: Species biometrics used in the CRM. Mean values and standard deviations (where applicable).

Species	Body length ¹ (m)	Wingspan ² (m)	Flight speed ³ (m/s)	Nocturnal Activity Rate ⁴ (%)	Avoidance rate ⁵	Flight type
Arctic tern	0.33 (0)	0.87 (0)	10.5 (0)	0	99.1 (0.04)	Flapping
Black-headed gull	0.37 (0)	1.1 (0)	11.9 (0)	25.0 (0)	99.5 (0.02)	Flapping
Common gull	0.42 (0)	1.3 (0)	13.4 (0)	25.0 (0)	99.5 (0.02)	Flapping
Common tern	0.33 (0)	0.87 (0)	10.5 (0)	0	99.1 (0.04)	Flapping
Fulmar	0.48 (0)	1.07 (0)	13.0 (0)	75 (0)	99.1 (0.04)	Flapping
Gannet	0.94 (0.0325)	1.72 (0.0375)	14.9 (0)	8 (10.0)	99.79 (0.03)	Flapping
Great black-backed gull	0.71 (0.035)	1.58 (0.0375)	13.7 (1.2)	37.5 (6.375)	99.4 (0.04)	Flapping
Herring gull	0.6 (0.0225)	1.44 (0.03)	12.8 (1.8)	37.5 (6.375)	99.4 (0.04)	Flapping
Kittiwake	0.39 (0.05)	1.08 (0.0625)	8.71 (0.4)	37.5 (6.375)	99.3 (0.03)	Flapping
Lesser black-backed gull	0.58 (0.03)	1.42 (0.0375)	13.1 (1.9)	37.5 (6.375)	99.4 (0.04)	Flapping
Little gull	0.26 (0)	0.78 (0)	12.2 (0)	25.0 (0)	99.5 (0.02)	Flapping
Sandwich tern	0.39 (0.005)	1.0 (0.04)	10.3 (3.4)	0 (0)	99.1 (0.04)	Flapping

1. Robinson (2005)

2. Pennycuik (1987), Alerstam (2007), Skov *et al.*, (2018)

3. Garthe and Hüppop (2004), Furness *et al.* (2018), MacArthur Green (2015)

4. Cook *et al.* (2014), JNCC *et al.*, (2014), Bowgen and Cook (2018)

5. Natural England (2022)

Table 12.4.2: Monthly mean densities (and 95% confidence intervals) of birds in flight within the Array Area used in the deterministic CRM.

Species	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Arctic tern	0 (0-0)	0 (0-0)	0 (0-0)	3.78 (0.61-9.8)	1.39 (0.39-3.06)	0 (0-0)	0 (0-0)	25.33 (11.11-41.47)	0.46 (0-1.16)	0.31 (0-0.74)	0 (0-0)	0 (0-0)
Black-headed gull	6.49 (1.85-13.72)	5.71 (1.08-12.87)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0.08 (0-0.27)	0.08 (0-0.23)	0 (0-0)	0.85 (0.15-1.85)	2.78 (0.65-5.64)	6.87 (1.04-18.44)
Common gull	9.73 (1.46-20.4)	26.41 (7.45-49.72)	0.08 (0-0.23)	0 (0-0)	0 (0-0)	0.46 (0-1.39)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	2.93 (0.93-6.37)	7.26 (0.42-20.2)
Common tern	0 (0-0)	0 (0-0)	0 (0-0)	1 (0.46-1.86)	0 (0-0)	0 (0-0)	0 (0-0)	11.74 (4.54-20.25)	1.7 (0.46-2.93)	0 (0-0)	0 (0-0)	0 (0-0)
Fulmar	0 (0-0)	0 (0-0)	0 (0-0)	0.08 (0-0.23)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0.15 (0-0.46)	0 (0-0)	0 (0-0)	0 (0-0)
Gannet	0.08 (0-0.23)	0 (0-0)	0.15 (0-0.46)	0 (0-0)	0 (0-0)	0.23 (0-0.74)	0.15 (0-0.39)	0.23 (0-0.58)	0.08 (0-0.27)	0.39 (0-0.93)	0 (0-0)	0.15 (0-0.46)
Great black-backed gull	0.15 (0-0.46)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Herring gull	0 (0-0)	0.08 (0-0.27)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0.08 (0-0.23)	0 (0-0)
Kittiwake	14.98 (6.09-24.08)	79 (33.19-147.95)	1.85 (0.42-4.37)	1.54 (0.31-3.24)	3.48 (1.27-6.88)	0.39 (0.04-0.97)	1.24 (0.18-2.57)	1.31 (0.58-2.21)	0.93 (0.23-1.89)	2.32 (0.7-4.37)	8.96 (3.43-16.66)	17.61 (3.61-41.61)
Lesser black-backed gull	0 (0-0)	0 (0-0)	0.08 (0-0.27)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
Little gull	7.72 (0.31-19.13)	1.31 (0.08-3.48)	3.4 (0-9.9)	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0.39 (0-0.93)	2.47 (0.08-6.03)	2.86 (0.15-6.95)	7.95 (1.97-15.96)	14.98 (0.89-29.62)
Sandwich tern	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)	0.08 (0-0.23)	0 (0-0)	0 (0-0)	0.15 (0-0.46)	0.23 (0-0.46)	0 (0-0)	0 (0-0)	0 (0-0)

Table 12.4.3: Windfarm and turbine specifications used in the CRM.

Turbine scenario	Average Revolutions Per Minute (RPM)	Rotor radius (m)	Hub height above Lowest Astronomical Tide (LAT) (m)	Max. blade width (m)	Mean blade pitch (°)	No. of turbines	Latitude (°)	Tidal offset (m, LAT to Mean Sea Level (MSL))
1a	6.34	118	155	5.4	10	56	52.81	1.115
1b	5.73	118	155	6.8	10	56	52.81	1.115
2	6.19	125	162	6.9	10	47	52.81	1.115

Table 12.4.4: Wind turbine Monthly percentage operational time used in the CRM.

Turbine scenario	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1a	0.958	0.994	0.959	0.921	0.906	0.916	0.863	0.918	0.931	0.98	0.959	0.969
1b	0.958	0.994	0.959	0.921	0.906	0.916	0.863	0.918	0.931	0.98	0.959	0.969
2	0.964	0.989	0.962	0.927	0.917	0.922	0.876	0.925	0.943	0.981	0.966	0.974

1.3 References

- Alerstam, T. *et al.* (2007). Flight speeds among bird species: Allometric and phylogenetic effects. *PLoS Biology*, 5(8), pp. 1656–1662. doi: 10.1371/journal.pbio.0050197.
- Band, W. (2012) Using a collision risk model to assess bird collision risks for offshore windfarms. The Crown Estate Strategic Ornithological Support Services (SOSS) report SOSS-02. SOSS Website.
- Bowgen, K. and Cook, A. (2018) Bird Collision Avoidance: Empirical evidence and impact assessments. JNCC Report No. 614, JNCC, Peterborough, ISSN 0963-8091.
- Cook, A.S.C.P., Humphries, E.M., Masden, E.A., and Burton, N.H.K. (2014). The avoidance rates of collision between birds and offshore turbines. BTO research Report No 656 to Marine Scotland Science.
- Furness, R.W., Garthe, S., Trinder, M., Matthiopoulos, J., Wanless, S. and Jeglinski, J. (2018). Nocturnal flight activity of northern gannets *Morus bassanus* and implications for modelling collision risk at offshore windfarms. *Environmental Impact Assessment Review*, 73, pp.1-6.
- Garthe, S and Hüppop, O. (2004) Scaling possible adverse effects of marine windfarms on seabirds: developing and applying a vulnerability index. *Journal of Applied Ecology*, 41, 724-734.
- Johnston, A., Cook, A.S.C.P., Wright, L.J., Humphreys, E.M. and Burton, E.H.K. (2014) Modelling flight heights of marine birds to more accurately assess collision risk with offshore wind turbines. *Journal of Applied Ecology*, 51, 31-41.
- JNCC, Natural England, NIEA, NRW, SNH (2014) Joint Response from the Statutory Nature Conservation Bodies to the Marine Scotland Science Avoidance Rate Review.
- Natural England (2022). Natural England interim advice on updated Collision Risk Modelling parameters (July 2022).
- Ozsanlav-Harris, L., Inger, R. & Sherley, R. (2022). Review of data used to calculate avoidance rates for collision risk modelling of seabirds. JNCC Report 732 (Research and review report), JNCC, Peterborough, ISSN 0963-8091. <https://hub.jncc.gov.uk/assets/de5903fe-81c5-4a37-a5bc-387cf704924d>.
- Pennyquick, C.J. (1987). Flight of auks (Alcidae) and other northern seabirds compared with southern Procellariiformes: ornithodolite observations. *Journal of Experimental Biology*, 128(1), pp.335-347.
- Robinson, R.A. (2005). BirdFacts: profiles of birds occurring in Britain & Ireland. BTO, Thetford (<http://www.bto.org/birdfacts>, accessed on 12 January 2021).
- Skov, H., Heinänen, S., Norman, T., Ward, R.M., Méndez-Roldán, S. and Ellis, I. 2018. ORJIP Bird Collision and Avoidance Study. Final report – April 2018. The Carbon Trust. 247 pp.